



# Mark Scheme (Results)

Summer 2019

Pearson Edexcel International GCSE  
In Mathematics A (4MA1)  
Paper 1H

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2019

Publications Code 4MA1\_1H\_1906\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2019

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.

Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
  - M marks: method marks
  - A marks: accuracy marks
  - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
  - cao – correct answer only
  - ft – follow through
  - isw – ignore subsequent working
  - SC - special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - awrt – answer which rounds to
  - eeo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods mark the one that leads to the answer on the answer line. If there is no answer given then mark the method that gives the lowest mark and award this mark.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

International GCSE Maths					
Apart from questions 1, 11, 12b, 15 (where the mark scheme states otherwise) the correct answer, unless clearly obtained from an incorrect method, should be taken to imply a correct method.					
Question	Working	Answer	Mark	Notes	
1	e.g. $\frac{14}{3}$ <b>and</b> $\frac{10}{9}$		3	M1	Both fractions expressed as improper fractions
	e.g. $\frac{14}{3} \times \frac{9}{10}$			M1	<b>or</b> for both fractions expressed as equivalent fractions with denominators that are a common multiple of 3 and 9 eg. $\frac{42}{9} \div \frac{10}{9}$ <b>or</b> $\frac{126}{27} \div \frac{30}{27}$
	e.g. $\frac{14}{3} \times \frac{9}{10} = \frac{126}{30} = \frac{21}{5} = 4\frac{1}{5}$ <b>or</b> $\frac{14}{3} \times \frac{9}{10} = \frac{126}{30} = 4\frac{6}{30} = 4\frac{1}{5}$ <b>or</b> $\frac{14^7}{3^1} \times \frac{9^3}{10^5} = \frac{21}{5} = 4\frac{1}{5}$ <b>or</b> $\frac{126}{27} \div \frac{30}{27} = \frac{126}{30} = \frac{21}{5} = 4\frac{1}{5}$	Shown		A1	Dep on M2 for conclusion to $4\frac{1}{5}$ from correct working – either sight of the result of the multiplication e.g. $\frac{126}{30}$ must be seen or correct cancelling prior to the multiplication to $\frac{21}{5}$ NB: use of decimals scores no marks
				Total 3 marks	

2	(a)	15 km/h or $\frac{25}{6}$ m/sec or 0.25 km/min or $\frac{15}{4}$ oe 12 km/h or $\frac{10}{3}$ m/sec or 0.2 km/min or $\frac{9}{3}$ oe	'before' with reason	1	B1 e.g. before as gradient is steeper <b>or</b> before as speed before is 15 km/h speed after is 12 km/h <b>or</b> before as she goes over 11(allow 11-12) km in $\frac{3}{4}$ hour but only goes 9 km in $\frac{3}{4}$ hour after oe NB: any figures used for the reason must be accurate if they haven't used 'gradient is steeper' oe
	(b)		line from (12:00, 24) to (12:45, 24) to (14:15, 0)	2	B2 If not B2 then B1 for a line from (12:00, 24) to (12:45, 24) <b>or</b> for a line from (t, 24) to (t + 1.5, 0) <b>or</b> for a time of 1.5 hours (oe) seen
	(c)	1h 45m + 1h 30m <b>or</b> 1 + 0.75 + 1.5 <b>or</b> 3h 15m <b>or</b> 3.25h <b>or</b> 195m oe		3	M1 ft from their graph for total time when cycling
		(24 × 2) ÷ "3.25" oe eg (48 ÷ 195) × 60			M1 ft dep on M1 for full method
			14.8		A1 awrt 14.8
				<b>Total 6 marks</b>	

<b>3</b>	(a)		$e^4$	1	B1
	(b)		$y^{16}$	1	B1
	(c)	$x^2 + 9x - 2x - 18$		2	M1 for 3 correct terms <b>or</b> 4 correct terms ignoring signs <b>or</b> $x^2 + 7x + c$ <b>or</b> .... + 7x - 18
			$x^2 + 7x - 18$		A1
	(d)		$4cp^2(4c^3 + 5p)$	2	B2 if not B2 then award B1 for any correct factorisation with at least 2 factors outside the bracket eg $4cp(4c^3p + 5p^2)$ , $cp^2(16c^3 + 20p)$ , $2p(8pc^4 + 10cp^2)$ etc <b>or</b> the correct common factor <b>and</b> a 2 term expression with just one error
					<b>Total 6 marks</b>

<b>4</b>	(a)		9, 3, (-1), -3, (-3), -1, (3)	2	B2 If not B2 then award B1 for at least 2 correct values
	(b)			2	M1 dep on B1 ft from (a) for at least 5 points plotted correctly
			correct graph		A1 for the correct graph (clear intention to go through all the points and which must be curved at the bottom)
					<b>Total 4 marks</b>

<b>5</b>		$2x + 0.18 + 2x + 3x + 0.26 + x = 1$ <b>or</b> $1 - (0.18 + 0.26) (= 0.56)$		4	M1
		$x = (1 - 0.18 - 0.26) \div (2 + 2 + 3 + 1) (=0.07)$			M1
		eg $(0.18 + 4 \times "0.07") \times 200$ <b>or</b> $0.46 \times 200$ <b>or</b> $36 + 42 + 14$ oe			M1 dep on M2 and probabilities between 0 and 1 <b>or</b> $\frac{92}{200}$ , oe with 92 seen
			92		A1
					<b>Total 4 marks</b>

<b>6</b>		$12 \times 8 \times 5 (= 480)$		3	M1
		"480" $\times 0.7$			M1 Dep on M1
			336		A1
					<b>Total 3 marks</b>



<b>7</b>	(a)		5 700 000	1	B1
	(b)		$4 \times 10^{-3}$	1	B1
	(c)		5 000 000 <b>or</b> $5 \times 10^6$ oe	2	B2 If not B2 then award B1 for 320000 or $3.2 \times 10^5$ oe or $5 \times 10^n$ oe where $n \neq 6$
					<b>Total 4 marks</b>

<b>8</b>		$0.08 \times 170\,000 (=13600)$ <b>or</b> $0.92 \times 170\,000 (=156400)$		3	M1 oe eg $170\,000 \div 12.5$	M2 for $170\,000 \times 0.92^3$
		e.g. $0.92 \times (0.92 \times "156400")$			M1 (dep)for a complete method	
			132377		A1 or 132376.96	
					(SCB2 for $170\,000 \times 0.92^4 (=121786.810)$ ) (SCB1 for $170\,000 \times 0.24 (=40\,800)$ <b>or</b> $170\,000 \times 0.76 (=129\,200)$ <b>or</b> $170\,000 \times 1.08 (=183\,600)$ <b>or</b> $170\,000 \times 1.08^3 (=214151)$ <b>or</b> an answer of $129\,200$ <b>or</b> an answer of $214\,151 - 214151.1(0)$ )	
					<b>Total 3 marks</b>	

9		$0.5 \times 6 \times 6 (=18)$		5	M1 For area of triangle, or may use $\frac{1}{2} \times 6 \times 6\sqrt{2} \sin 45$ or $\frac{1}{2} \times 6\sqrt{2} \times 3\sqrt{2}$ oe
		$(d^2 =) 6^2 + 6^2 (=72)$ or $\frac{AC}{(\sin 90)} = \frac{6}{\sin 45}$			M1
		$\sqrt{6^2 + 6^2} (= \sqrt{72} = 6\sqrt{2} = 8.4(85\dots)\text{or } 8.5)$ <b>or</b> $AC = \frac{6(\sin 90)}{\sin 45} = 6\sqrt{2} = 8.4(85\dots)\text{or } 8.5$ oe			M1
		$0.5 \times \pi \times \left(\frac{"8.48.."}{2}\right)^2 (= 9\pi \text{ or } 28\dots)$			M1
			46.3		A1 for 46.2 – 46.3
					<b>Total 5 marks</b>

10		$(8 =) 2 \times 2 \times 2$ or $2^3$ or $2^{3+n}$		2	M1 For clearly writing 8 as a product of prime factors or as $2^3$
			$2^{n+3} \times 3 \times 5^m$		A1
					<b>Total 2 marks</b>

11		5.5 or 6.5 or 12.5 or 17.5		3	M1 Accept 6.49 for 6.5 and 17.49 for 17.5
		17.5 – 5.5			M1 for UB – LB where $15 < UB \leq 17.5$ and $5.5 \leq LB < 6$
			12		A1 dep on M2
					<b>Total 3 marks</b>

12	(a)		$(2x - 3)(x - 2)$	2	B2 or $(3 - 2x)(2 - x)$ (B1 for $(2x + a)(x + b)$ where $ab = 6$ or $2b + a = -7$ eg $(2x + 3)(x + 2)$ , $(2x - 5)(x - 1)$ ), etc or for
	(b)	$4m + 9 = 3(7 - 2m)$		4	M1 for removing fraction
		$4m + 9 = 21 - 6m$			M1 for correct expansion of bracket in a correct equation
		$4m + 6m = 21 - 9$ <b>or</b> $10m = 12$ <b>or</b> $-21 + 9 = -6m - 4m$ <b>or</b> $-10m = -12$			M1 for a correct equation with $m$ terms isolated on one side ft their equation if first M1 awarded
			$\frac{12}{10}$ oe		A1 dep on at least M2 [SC: B2 for an answer of $m = 2$ with working shown (from $4m + 9 = 21 - 2m$ oe) <b>or</b> $m = -0.2$ oe with working shown (from $4m + 9 = 7 - 6m$ oe)]
		<b>Alternative</b>			
		$\frac{4}{3}m + 3 = 7 - 2m$		4	M1 Division of each term on LHS by 3
		$\frac{4}{3}m + 2m = 7 - 3$ oe			M1 for a correct equation with $m$ terms isolated on one side ft their equation if first M1 awarded

		$10m = 3 \times 4$ oe			M1 For removing fraction in a fully correct equation
			$\frac{12}{10}$ oe		A1 dep on at least M2
<b>12 contd</b>	(c)	$\frac{y^{\frac{1}{4}}}{y}$ <b>or</b> $\sqrt[4]{y} = y^{\frac{1}{4}}$ <b>or</b> $y^{\frac{1}{4}-1}$		2	M1 or $b = -\frac{3}{4}$
			$y^{-\frac{3}{4}}$		A1
					<b>Total 8 marks</b>

13	(a)		$\frac{6}{14}, \frac{8}{14}$	2	B1 for $\frac{6}{14}\left(\frac{3}{7}\right), \frac{8}{14}\left(\frac{4}{7}\right)$ in correct positions. Allow decimals of 2dp or better (0.43, 0.57)
			$\frac{3}{10}, \frac{7}{10}, \frac{3}{10}, \frac{7}{10}$		B1oe for $\frac{3}{10}, \frac{7}{10}, \frac{3}{10}, \frac{7}{10}$ in correct positions.
	(b)	$\frac{8}{14} \times \frac{7}{10}$		2	M1 ft from (a)
			$\frac{2}{5}$ oe		A1
	(c)	$\frac{7}{13} \times \frac{6}{9} \left( = \frac{42}{117} = \frac{14}{39} = 0.35(897...) \right)$ <b>or</b> $\frac{8}{14} \times \frac{7}{13} \left( = \frac{56}{182} \text{ oe} \right)$ <b>or</b> $\frac{7}{10} \times \frac{6}{9} \left( = \frac{42}{90} \right)$		3	M1 ft from (a) $\left( \frac{7}{13} = 0.54 \text{ to 2dp} \right)$ $\frac{6}{9} = 0.67 \text{ to 2dp}$
		" $\frac{42}{117}$ " " $\times$ " " $\frac{2}{5}$ " or $\left( \frac{8}{14} \times \frac{7}{13} \right) \times \left( \frac{7}{10} \times \frac{6}{9} \right)$			M1 ft from (b)
			$\frac{28}{195}$ oe		A1 for $\frac{28}{195}$ oe, e.g. 0.14(3589...) from accurate working
					<b>Total 7 marks</b>

<b>14</b>	(a)		7, 8, 9, 10, 11	2	B2 completely correct. (B1 for 4 or 5 correct and no more than 1 incorrect <b>or</b> for all terms seen correctly placed in a Venn diagram or for a correct description of the numbers in the set but not listed, eg $7 \leq x < 12$ )
	(b)		eg 2, 4, 6	1	B1 for any 3 of 2, 4, 6, 8, 10
					<b>Total 3 marks</b>

15		$x = 0.25454\ldots$ $100x = 25.454\ldots$ $10x = 2.5454\ldots$ $1000x = 254.54\ldots$		2	<p>M1 For 2 recurring decimals that when subtracted give a whole number or terminating decimal eg 25.2 or 252 etc eg <math>100x = 25.454\ldots</math> and <math>x = 0.25454\ldots</math> <b>or</b> <math>1000x = 254.54\ldots</math> and <math>10x = 2.5454\ldots</math> with intention to subtract.          (if recurring dots not shown then showing at least the digits 25454, ie 5sf)  <b>or</b>  <math>0.2 + 0.0\dot{5}4</math> <b>and</b>          eg <math>x = 0.05454\ldots</math>, <math>100x = 5.4545\ldots</math> with intention to subtract.</p>
		<p>eg <math>100x - x = 25.454\ldots - 0.254\ldots = 25.2</math> <b>and</b>  <math>\frac{25.2}{99} = \frac{14}{55}</math> <b>or</b>  <math>1000x - 10x = 254.545\ldots - 2.545\ldots = 252</math> <b>and</b>  <math>\frac{252}{990} = \frac{14}{55}</math> <b>or</b></p> <p><math>100x - x = 5.4545\ldots - 0.05454\ldots = 5.4</math> <b>and</b>  <math>\frac{5.4}{99} = \frac{54}{990} \left( = \frac{3}{55} \right)</math> <b>and</b> <math>\frac{2 \times 99 + 54}{990} = \frac{252}{990} = \frac{14}{55}</math>          or <math>\frac{5.4}{99} = \frac{54}{990} = \frac{3}{55}</math> and <math>\frac{11+3}{55} = \frac{14}{55}</math></p>	show		<p>A1 for completion to <math>\frac{14}{55}</math></p>



					<b>Total 2 marks</b>
<b>16</b>		$a = 7$ and $d = 3$ $\frac{100}{2}(2 \times 7 + (100 - 1) \times 3)$ <b>or</b> 100th term is $7 + (100 - 1) \times 3 (= 304)$ <b>and</b> $100 \times (7 + "304") \div 2$ <b>or</b>  100 <sup>th</sup> term is $3 \times 100 + 4 (= 304)$ <b>and</b> $100 \times (7 + "304") \div 2$		2	M1 for a method to find the sum - brackets $(100 - 1)$ must be used correctly
			15 550		A1
					<b>Total 2 marks</b>

17	(a)	eg $\frac{24}{36}$ or 2 : 3 oe <b>or</b> $\frac{36}{24}$ or 3 : 2 oe		2	M1 for a correct scale factor
			2160		A1
	(b)	$\left(\frac{24}{36}\right)^3$ or $2^3 : 3^3$ oe <b>or</b> $\left(\frac{36}{24}\right)^3$ or $3^3 : 2^3$ oe or $\frac{8}{27}$ or $\frac{27}{8}$ oe		2	M1 For correct SF for volume ft from linear scale factor in (a) or ft from $\sqrt{\frac{2160}{960}}$
					$(A =) \frac{8}{27} V$ oe
					<b>Total 4 marks</b>

18		$17.8^2 + 26.3^2 - 2 \times 17.8 \times 26.3 \times \cos 36$		3	M1
		e.g. 1008.5... – 757.... <b>or</b> 251(.06...)			M1 for correct order of operations
			15.8		A1 for ans in range 15.8 – 15.9
					<b>Total 3 marks</b>

19		$15 \div 20 (=0.75)$ $48 \div 15 (=3.2)$ $21 \div 5 (=4.2)$ $16 \div 10 (=1.6)$	correct histogram	3	B3 For a fully correct histogram [If not B3 then B2 for 3 correct frequency densities (can be implied by heights) or 3 correct bars drawn If not B2 then B1 for 2 correctly calculated frequency densities (can be implied by heights) or 2 correct bars drawn.]
				Total 3 marks	

Students can use other methods to gain the correct answer					
20		angle $ABD = 71$ <b>or</b> angle $ACD = 71$ <b>or</b> using $O$ as centre of circle, angle $ADO = 90 - 71 (=19)$		5	M1 clearly labelled or stated
		angle $ADB = 71$ <b>or</b> angle $ACB = 71$ <b>or</b> angle $BAD = 19 \times 2 (=38)$ <b>or</b> reflex angle $BOD = 2 \times 142 (=284)$			M1 dep clearly labelled or stated
		angle $BCD = 142$	142		A1 Clearly labelled or stated, from no incorrect working for their method
					B2 dep on A1 for fully correct reasons for each stage of working, repeated if used more than once. eg <u>alternate segment</u> theorem, base angles in an <u>isosceles</u> triangle are equal, <u>angles</u> in a <u>triangle</u> sum to $180^\circ$ , angle between <u>tangent</u> and <u>radius(diameter)</u> is $90^\circ$ <u>congruent</u> triangles ( <u>equal</u> triangles) oe opposite angles of a <u>cyclic quadrilateral</u> sum to $180^\circ$ <u>angles</u> in the <u>same segment</u> <u>angle</u> at the <u>centre</u> is $2 \times$ angle at <u>circumference</u> oe <u>equal chords</u> subtend <u>equal angles</u> at the <u>circumference</u> If not B2 then award B1 dep on M1 for any one correct circle theorem reason associated with angle(s) found
					<b>Total 5 marks</b>


21		$h = 3r$ or $r = \frac{h}{3}$		5	M1	for $h = 3r$ <b>or</b> $r = \frac{h}{3}$ oe stated or used correctly
		$\frac{1}{2} \times \frac{4}{3} \times \pi r^3$ oe <b>or</b> $\pi \times r^2 \times 3r$ oe			M1	<b>or</b> $\frac{1}{2} \times \frac{4}{3} \pi \left(\frac{h}{3}\right)^3$ or $\pi \left(\frac{h}{3}\right)^2 h$
		$\frac{1}{2} \times \frac{4}{3} \times \pi r^3 + \pi \times r^2 \times 3r = 792\pi$ oe			M1	<b>or</b> $\frac{1}{2} \times \frac{4}{3} \pi \left(\frac{h}{3}\right)^3 + \pi \left(\frac{h}{3}\right)^2 h = 792\pi$
		$(r = ) 6$ or $(h = ) 18$			A1	
			24		A1ft	their " $6$ " $\times 4$ <b>or</b> " $18$ " $\times \frac{4}{3}$ correctly evaluated dep on M3
					<b>Total 5 marks</b>	

22	(a)		correct graph (see end of mark scheme) [must go through (60, 2), (150, 0), (240, -2), (330, 0)] and not through (0, 0)	2	B2 if not B2 then award B1 for a graph of the correct shape going through 2 or 3 of the given points or for a clear stretch of SF2 (ie a maximum point on graph at $(x_1, 2)$ and a minimum point at $(x_2, -2)$ ) or a clear translation of $\begin{pmatrix} -30 \\ 0 \end{pmatrix}$ (ie a point on graph at $(150, y)$ and a point at $(330, y)$ )
	(b)(i)		$(x - 3)^2 + 1$	2	B2 (B1 for $(x - \frac{6}{2})^2 + n$ (where $n \neq 1$ ) <b>or</b> for $(x - m)^2 + 1$ (where $m \neq 3$ ) <b>or</b> for $x^2 - ax - ax + a^2 + b$ with $2a = 6$ or $a^2 + b = 10$ )
	(b)(ii)		translation of $\begin{pmatrix} 3 \\ 1 \end{pmatrix}$	2	B1 for translation
					B1 For $\begin{pmatrix} 3 \\ 1 \end{pmatrix}$ ft from (b)(i) must be column vector
					<b>Total 6 marks</b>

<b>23</b>		$\left(\frac{10+2}{2}, \frac{7+19}{2}\right)$ <b>or</b> (6, 13)		5	M1
		$\frac{19-7}{10-2}\left(=\frac{12}{8}\right)$ oe <b>or</b> 1.5 oe			M1
		$m \times \frac{3}{2} = -1$ oe <b>or</b> $m = -\frac{2}{3}$			M1 for use of $m_1m_2 = -1$
		"13" = $-\frac{2}{3}$ "x "6" + c <b>or</b> $c = 17$ oe <b>or</b> $y - "13" = -\frac{2}{3}(x - "6")$			M1 Or for $y = -\frac{2}{3}x + 17$ [NB: "13", "6" and $-\frac{2}{3}$ " must come from correct working]
			$3y + 2x = 51$		A1 for $3y + 2x = 51$ <b>or</b> $3y = -2x + 51$ etc but must be integer coefficients
					<b>Total 5 marks</b>

<b>24</b>		$(v =) 3t^2 - 6 \times 2t + 5 (+ 0)$		4	M1 for differentiating at least 2 terms correctly
		$(a =) 3 \times 2t - 12$			M1 dep ft
		$6t - 12 = 3$			M1 dep on at least M1 for equating their acceleration in terms of $t$ to 3
			2.5 oe		A1
					<b>Total 4 marks</b>

**Q19**





q22

