

GCE Examinations

Further Pure Mathematics Module FP3

Advanced Subsidiary / Advanced Level

Paper A

Time: 1 hour 30 minutes

Instructions and Information

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 8 questions.

Advice to Candidates

You must show sufficient working to make your methods clear to an examiner.
Answers without working will gain no credit.



Written by Rosemary Smith & Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

1. With respect to a fixed origin O , the lines l_1 and l_2 are given by the equations

$$l_1 : [\mathbf{r} - (-3\mathbf{i} + 2\mathbf{j} - \mathbf{k})] \times (\mathbf{i} + \mathbf{k}) = 0,$$

$$l_2 : [\mathbf{r} - (\mathbf{i} + \mathbf{j} + 4\mathbf{k})] \times (2\mathbf{i} - \mathbf{j} - 2\mathbf{k}) = 0.$$

(a) Find $(\mathbf{i} + \mathbf{k}) \times (2\mathbf{i} - \mathbf{j} - 2\mathbf{k})$. **(3 marks)**

(b) Find the shortest distance between l_1 and l_2 . **(3 marks)**

2. Prove by induction that, for all $n \in \mathbb{Z}^+$,

$$\sum_{r=1}^n (r^2 + 1)r! = n(n+1)! \quad \text{(6 marks)}$$

3. (a) Solve the equation

$$z^3 + 27 = 0,$$

giving your answers in the form $re^{i\theta}$ where $r > 0$, $-\pi < \theta \leq \pi$. **(5 marks)**

(b) Show the points representing your solutions on an Argand diagram. **(2 marks)**

4.
$$\mathbf{A} = \begin{pmatrix} 2 & a \\ 2 & b \end{pmatrix}.$$

The matrix \mathbf{A} has eigenvalues $\lambda_1 = -2$ and $\lambda_2 = 3$.

(a) Find the value of a and the value of b . **(4 marks)**

Using your values of a and b ,

(b) for each eigenvalue, find a corresponding eigenvector, **(3 marks)**

(c) find a matrix \mathbf{P} such that $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} -2 & 0 \\ 0 & 3 \end{pmatrix}$. **(2 marks)**

5. $(1 + x^2) \frac{d^2y}{dx^2} + 4x \frac{dy}{dx} + 2y = 0$ and $y = 1, \frac{dy}{dx} = 1$ at $x = -1$.

Find a series solution of the differential equation in ascending powers of $(x + 1)$ up to and including the term in $(x + 1)^4$.

(11 marks)

6. The variable y satisfies the differential equation

$$\frac{d^2y}{dx^2} = x \frac{dy}{dx} + y^2 \quad \text{with } y = 1.2 \text{ at } x = 0.1 \quad \text{and } y = 0.9 \text{ at } x = 0.2$$

Use the approximations $\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_{-1}}{2h}$ and $\left(\frac{d^2y}{dx^2}\right)_0 \approx \frac{y_1 - 2y_0 + y_{-1}}{h^2}$ with a step length of 0.1 to estimate the values of y at $x = 0.3$ and $x = 0.4$ giving your answers to 3 significant figures.

(11 marks)

7.
$$\mathbf{M} = \begin{pmatrix} 2 & 1 & 1 \\ k & 4 & 3 \\ -1 & k & 2 \end{pmatrix}.$$

(a) Find the determinant of \mathbf{M} in terms of k . **(2 marks)**

(b) Prove that \mathbf{M} is non-singular for all real values of k . **(2 marks)**

(c) Given that $k = 3$, find \mathbf{M}^{-1} , showing each step of your working. **(4 marks)**

When $k = 3$ the image of the vector $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$ when transformed by \mathbf{M} is the vector $\begin{pmatrix} 0 \\ 3 \\ 5 \end{pmatrix}$.

(d) Find the values of a, b and c . **(3 marks)**

Turn over

8. A transformation T from the z -plane to the w -plane is defined by

$$w = \frac{z+1}{iz-1}, \quad z \neq -i,$$

where $z = x + iy$, $w = u + iv$ and x, y, u and v are real.

T transforms the circle $|z| = 1$ in the z -plane onto a straight line L in the w -plane.

(a) Find an equation of L giving your answer in terms of u and v . **(5 marks)**

(b) Show that T transforms the line $\text{Im } z = 0$ in the z -plane onto a circle C in the w -plane, giving the centre and radius of this circle. **(6 marks)**

(c) On a single Argand diagram sketch L and C . **(3 marks)**

END