

GCE Examinations

Further Pure Mathematics

Module FP3

Advanced Subsidiary / Advanced Level

Paper B

Time: 1 hour 30 minutes

Instructions and Information

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 8 questions.

Advice to Candidates

You must show sufficient working to make your methods clear to an examiner.
Answers without working will gain no credit.



Written by Rosemary Smith & Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

1. Given that x is so small that terms in x^3 and higher powers of x may be neglected, find the values of the constants a and b for which

$$\frac{\ln(1+ax)}{1+bx} = 3x + \frac{3}{2}x^2. \quad \text{(5 marks)}$$

2. Given that

$$|z + 1 - 4i| = 1,$$

(a) sketch, in an Argand diagram, the locus of z , (2 marks)

(b) find the maximum value of $\arg z$ in degrees to one decimal place. (3 marks)

3. (a) Show that

$$\cosh ix = \cos x \quad \text{where } x \in \mathbb{R}. \quad \text{(2 marks)}$$

(b) Hence, or otherwise, solve the equation

$$\cosh ix = e^{ix}$$

for $0 \leq x < 2\pi$. (3 marks)

4. Given that

$$u_{n+2} = 5u_{n+1} - 6u_n \quad \text{for } n \geq 1, \quad u_1 = 2 \text{ and } u_2 = 4,$$

prove by induction that $u_n = 2^n$ for all integers n , $n \geq 1$. (6 marks)

5.
$$\mathbf{M} = \begin{pmatrix} 1 & 2 & -1 \\ 0 & 1 & -4 \\ x & 3 & -1 \end{pmatrix}.$$

- (a) Given that $\lambda = -1$ is an eigenvalue of \mathbf{M} , find the value of x . **(3 marks)**
- (b) Show that $\lambda = -1$ is the only real eigenvalue of \mathbf{M} . **(6 marks)**
- (c) Find an eigenvector corresponding to the eigenvalue $\lambda = -1$. **(2 marks)**
-

6. A student is looking at different methods of solving the differential equation

$$\frac{dy}{dx} = xy \quad \text{with } y = 1 \text{ at } x = 0.2$$

The first method the student tries is to use the approximation $\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_0}{h}$ twice with a step length of 0.1 to obtain an estimate for y at $x = 0.4$

- (a) Find the value of the student's estimate for y at $x = 0.4$ **(6 marks)**

The student then realises that the exact value of y at $x = 0.4$ can be found using integration.

- (b) Use integration to find the exact value of y at $x = 0.4$ **(4 marks)**
- (c) Find, correct to 1 decimal place, the percentage error in the estimated value in part (a). **(2 marks)**
-

Turn over

7. (a) Given that $z = \cos\theta + i \sin\theta$, show that

$$z^n + \frac{1}{z^n} = 2 \cos n\theta \quad \text{and} \quad z^n - \frac{1}{z^n} = 2i \sin n\theta,$$

where n is a positive integer. **(3 marks)**

- (b) Given that

$$\cos^4\theta + \sin^4\theta = A \cos 4\theta + B,$$

find the values of the constants A and B . **(8 marks)**

- (c) Hence find the exact value of

$$\int_0^{\frac{\pi}{8}} \cos^4\theta + \sin^4\theta \, d\theta. \quad \text{(3 marks)}$$

8. The points A , B , C and D have coordinates $(3, -1, 2)$, $(-2, 0, -1)$, $(1, 2, 6)$ and $(-1, -5, 8)$ respectively, relative to the origin O .

- (a) Find $\overrightarrow{AB} \times \overrightarrow{AC}$. **(5 marks)**

- (b) Find the volume of the tetrahedron $ABCD$. **(3 marks)**

The plane Π contains the points A , B and C .

- (c) Find a vector equation of Π in the form $\mathbf{r} \cdot \mathbf{n} = p$. **(3 marks)**

The perpendicular from D to Π meets the plane at the point E .

- (d) Find the coordinates of E . **(6 marks)**

END