

GCE Examinations
Advanced / Advanced Subsidiary

Core Mathematics C4

Paper G

Time: 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures, unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphic calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**



Written by Shaun Armstrong

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

1. Express

$$\frac{2x^3 + x^2}{x^2 - 4} \times \frac{x - 2}{2x^2 - 5x - 3}$$

as a single fraction in its simplest form. [4]

2. A curve has the equation

$$x^3 + 2xy - y^2 + 24 = 0.$$

Show that the normal to the curve at the point $(2, -4)$ has the equation $y = 3x - 10$. [7]

3. Using the substitution $u = e^x - 1$, show that

$$\int_{\ln 2}^{\ln 5} \frac{e^{2x}}{\sqrt{e^x - 1}} dx = \frac{20}{3}. [8]$$

4. (i) Expand $(1 + ax)^{-3}$, $|ax| < 1$, in ascending powers of x up to and including the term in x^3 . Give each coefficient as simply as possible in terms of the constant a . [3]

Given that the coefficient of x^2 in the expansion of $\frac{6-x}{(1+ax)^3}$, $|ax| < 1$, is 3,

(ii) find the two possible values of a . [4]

Given also that $a < 0$,

(iii) show that the coefficient of x^3 in the expansion of $\frac{6-x}{(1+ax)^3}$ is $\frac{14}{9}$. [2]

5. $f(x) = \frac{7 + 3x + 2x^2}{(1-2x)(1+x)^2}$, $|x| > \frac{1}{2}$.

(i) Express $f(x)$ in partial fractions. [5]

(ii) Show that

$$\int_1^2 f(x) dx = p - \ln q,$$

where p is rational and q is an integer. [5]

6. Relative to a fixed origin, the points A , B and C have position vectors $(2\mathbf{i} - \mathbf{j} + 6\mathbf{k})$, $(5\mathbf{i} - 4\mathbf{j})$ and $(7\mathbf{i} - 6\mathbf{j} - 4\mathbf{k})$ respectively.

(i) Show that A , B and C all lie on a single straight line. [3]

(ii) Write down the ratio $AB : BC$ [1]

The point D has position vector $(3\mathbf{i} + \mathbf{j} + 4\mathbf{k})$.

(iii) Show that AD is perpendicular to BD . [4]

(iv) Find the exact area of triangle ABD . [3]

7. A mathematician is selling goods at a car boot sale. She believes that the rate at which she makes sales depends on the length of time since the start of the sale, t hours, and the total value of sales she has made up to that time, $\pounds x$.

She uses the model

$$\frac{dx}{dt} = \frac{k(5-t)}{x},$$

where k is a constant.

Given that after two hours she has made sales of $\pounds 96$ in total,


(i) solve the differential equation and show that she made $\pounds 72$ in the first hour of the sale. [7]

The mathematician believes that it is not worth staying at the sale once she is making sales at a rate of less than $\pounds 10$ per hour.

(ii) Verify that at 3 hours and 5 minutes after the start of the sale, she should have already left. [4]

Turn over

8.



The diagram shows the curve $y = f(x)$ in the interval $0 \leq x \leq 2\pi$ where

$$f(x) = \frac{\cos x}{2 - \sin x}, \quad x \in \mathbb{R}.$$

- (i) Show that $f'(x) = \frac{1 - 2 \sin x}{(2 - \sin x)^2}$. [3]
- (ii) Find an equation for the tangent to the curve $y = f(x)$ at the point where $x = \pi$. [3]
- (iii) Find the minimum and maximum values of $f(x)$ in the interval $0 \leq x \leq 2\pi$. [4]
- (iv) Explain why your answers to part (c) are the minimum and maximum values of $f(x)$ for all real values of x . [2]