



Pearson  
Edexcel

## Mark Scheme (Results)

Summer 2022

Pearson Edexcel GCE

A Level Further Mathematics (9FM0)

Paper 3B -Further Statistics 1

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2022

Question Paper Log Number P69194A\*

Publications Code 9FM0\_3B\_2206\_MS\*

All the material in this publication is copyright

© Pearson Education Ltd 2022

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

**EDEXCEL GCE MATHEMATICS**  
**General Instructions for Marking**

1. The total number of marks for the paper is 80.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - The second mark is dependent on gaining the first mark
4. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
  5. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response.  
If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete.
  6. Ignore wrong working or incorrect statements following a correct answer.

7. Mark schemes will firstly show the solution judged to be the most common response expected from candidates. Where appropriate, alternative answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used.

Qu	Scheme	Marks	AOs												
1(a)	$r = P(X = 3) \times 100$ or $r = P(X = 1) \times 100$ or $s = P(X = 2) \times 100$ $r = \underline{25}$ (value may be in table) $s = \underline{37.5}$ (value may be in table)	M1 A1 A1 <b>(3)</b>	3.4 1.1b 1.1b												
(b)	Ho: B(4,0.5) is a suitable model (o.e.) Condone B(0.5, 4) Hi: B(4,0.5) is not a suitable model (o.e.)	B1	2.5												
	<table border="1"> <tr> <td><math>\frac{(O_i - E_i)^2}{E_i}</math></td> <td>2.25</td> <td>2.56</td> <td>0.54</td> <td>4</td> <td>1.21</td> </tr> <tr> <td><math>\frac{O_i^2}{E_i}</math></td> <td>16</td> <td>43.56</td> <td>29.04</td> <td>9</td> <td>12.96</td> </tr> </table>	$\frac{(O_i - E_i)^2}{E_i}$	2.25	2.56	0.54	4	1.21	$\frac{O_i^2}{E_i}$	16	43.56	29.04	9	12.96	M1	1.1b
$\frac{(O_i - E_i)^2}{E_i}$	2.25	2.56	0.54	4	1.21										
$\frac{O_i^2}{E_i}$	16	43.56	29.04	9	12.96										
	$\sum \frac{(O_i - E_i)^2}{E_i} = 10.56$ or $\sum \frac{O_i^2}{E_i} - N = 110.56 - 100 = 10.56 \left( = \frac{264}{25} \right)$ $\nu = 5 - 1 = 4$ CV = 9.488 (Calc 9.487729035...)	A1 B1 B1ft	1.1b 1.1b 1.1b												
	Significant so there is evidence that the researcher's <b>model is not suitable</b>	A1	2.2b												
		<b>(6)</b>													

**Total 9**

(a)	<b>M1</b> Using the Binomial model to expected value. Allow if <u>both</u> probs 0.25 and 0.375 seen <b>1<sup>st</sup>A1</b> May be implied by a correct value of $r$ or $s$ . Alternatives $r = 6.25 \times 4$ or $s = 6.25 \times 6$ <b>2<sup>nd</sup>A1</b> for $r = 25$ for $s = 37.5$		
SC	<b>“B1”</b> If M0 scored but their values of $r$ and $s$ satisfy $2r + s = 87.5$ score as <b>M0A0A1</b>		
(b)	<b>1<sup>st</sup> B1</b> Both hypotheses correct using the correct notation in at least one <u>or</u> written in full e.g. binomial with $n = 4$ and $p = 0.5$ <b>M1</b> Calculating either $\frac{(O_i - E_i)^2}{E_i}$ or $\frac{O_i^2}{E_i}$ at least 4 correct. Implied by sight of awrt 10.6 <b>1<sup>st</sup> A1</b> Allow 10.6 (from correct working) <b>2<sup>nd</sup> B1</b> Correct dof May be implied by CV of 9.48 or 9.49 or better <b>3<sup>rd</sup>B1ft</b> For 9.488 or better. Can fit their dof NB $\chi_3^2(5\%) = 7.815$ (allow awrt 7.815) <b>2<sup>nd</sup>A1</b> <b>Indep’ of hypotheses but dep on 1<sup>st</sup> A1</b> Evaluating the outcome by drawing a correct inference. Compatible with comparison of 10.56 or 10.6 with their CV (which must be $> 1$ ) They must say <b>model not suitable</b> (o.e.) They do not need to state the comparison or say reject $H_0$ etc No need to explicitly see B(4, 0.5) mentioned here		

Question	Scheme	Marks	AOs
----------	--------	-------	-----

<b>2(a)</b>	$[E(X) = ]0.2b - 1$	B1 <b>(1)</b>	1.1b																																										
<b>(b)</b>	$E(X^2) = 25 \times 0.3 + 1 \times 0.25 [+0 \times 0.1] + 25 \times 0.15 + 0.2b^2 [= 11.5 + 0.2b^2]$ $"11.5 + 0.2b^2" - ("0.2b - 1")^2 [= 34.26]$ $0.16b^2 + 0.4b - 23.76 [= 0] \quad \text{or} \quad \frac{4}{25}b^2 + \frac{2}{5}b - \frac{594}{25} [= 0]$ $b = \underline{11}$ [since $b > 5$ ]	M1 M1 M1 A1 <b>(4)</b>	1.1b 3.1a 1.1b 2.2a																																										
<b>(c)</b>	<table border="1" style="margin-left: auto; margin-right: auto;"> <tbody> <tr><td><math>X</math></td><td>-5</td><td>-1</td><td>0</td><td>5</td><td>"11"</td></tr> <tr><td><math>X^2</math></td><td>25</td><td>1</td><td>0</td><td>25</td><td>"121"</td></tr> <tr><td><math>2 - 3X</math></td><td>17</td><td>5</td><td>2</td><td>-13</td><td>"-31"</td></tr> <tr><td><math>X^2 - 2</math></td><td>23</td><td>-1</td><td>-2</td><td>23</td><td>"119"</td></tr> <tr><td><math>-3X</math></td><td>15</td><td>3</td><td>0</td><td>15</td><td>"-33"</td></tr> <tr><td><math>X^2 + 3X</math></td><td>10</td><td>-2</td><td>0</td><td>40</td><td>"154"</td></tr> <tr><td><math>X^2 + 3X - 2</math></td><td>8</td><td>-4</td><td>-2</td><td>38</td><td>"152"</td></tr> </tbody> </table> $P(X^2 < 2 - 3X) = P(X = -1) + P(X = 0)$ $= \underline{0.35}$	$X$	-5	-1	0	5	"11"	$X^2$	25	1	0	25	"121"	$2 - 3X$	17	5	2	-13	"-31"	$X^2 - 2$	23	-1	-2	23	"119"	$-3X$	15	3	0	15	"-33"	$X^2 + 3X$	10	-2	0	40	"154"	$X^2 + 3X - 2$	8	-4	-2	38	"152"	M1 A1ft  M1 A1 <b>(4)</b>	2.1 1.1b  2.2a 1.1b
$X$	-5	-1	0	5	"11"																																								
$X^2$	25	1	0	25	"121"																																								
$2 - 3X$	17	5	2	-13	"-31"																																								
$X^2 - 2$	23	-1	-2	23	"119"																																								
$-3X$	15	3	0	15	"-33"																																								
$X^2 + 3X$	10	-2	0	40	"154"																																								
$X^2 + 3X - 2$	8	-4	-2	38	"152"																																								
<b>Total 9</b>																																													
<b>(a)</b>	<b>B1</b>	Correct expression for $E(X)$																																											
<b>(b)</b>	<b>1<sup>st</sup> M1</b>	Correct attempt at $E(X^2)$ using $\sum x^2 P(X = x)$ at least 3 correct non-zero products Allow $(-5)^2$ etc																																											
	<b>2<sup>nd</sup> M1</b>	Realising that $\text{Var}(X) = E(X^2) - [E(X)]^2$ needs to be <b>used</b>																																											
	<b>3<sup>rd</sup> M1</b>	Reducing their equation to a 3 term quadratic. At least 2 terms correct. Allow e.g. $0.16b^2 + 0.4b = 23.76$ Condone missing "=0"																																											
	<b>A1</b>	For 11 only (from the correct equation) so -13.5 must be eliminated Correct answer with no incorrect working seen scores 4/4																																											
<b>(c)</b>	<b>1<sup>st</sup> M1</b>	At least 4 values correct for $(X^2 \text{ and } 2 - 3X)$ <u>or</u> for $(X^2 - 2 \text{ and } -3X)$ <u>or</u> $X^2 + 3X$ <u>or</u> $X^2 + 3X - 2$ (o.e.) Allow for solving equation with one sign error																																											
	<b>1<sup>st</sup> A1ft</b>	All correct or correct ft with their $b$ but must have $b > 5$ (accurate to 1 sf) Allow solving equation to get awrt -3.6 <b>and</b> awrt 0.56 <u>or</u> $\frac{-3 \pm \sqrt{17}}{2}$ (ft their $b > 5$ )  If there are omissions <b>but no errors</b> in the lists of values then if 2 <sup>nd</sup> M1 and 2 <sup>nd</sup> A1 are scored then the 1 <sup>st</sup> M1 and 1 <sup>st</sup> A1 can be given by implication.																																											
	<b>2<sup>nd</sup> M1</b>	For identifying the correct values of $X$ required i.e. $X = -1$ and $X = 0$																																											
	<b>2<sup>nd</sup> A1</b>	0.35 <b>NB</b> It is possible to score M0A0M1A1 here if their table of values is incorrect Correct answer with no incorrect working seen scores 4/4 (Allow correct use of their $b > 5$ )																																											

Qu	Scheme	Marks	AOs
3(a)	$W \sim \text{Po}(11.2)$ and $P(W \leq 19) = 1 - P(W \geq 18)$ <u>or</u> suitable 3sf probs $P(W \leq 19) = 0.020776\dots$ awrt <b>0.021</b>	M1	3.4
		A1	1.1b
		(2)	
(b)	$[S = \# \text{ calls per day, } S \sim \text{Po}(0.4)]$ $P(S > 1) = 0.061551\dots$ awrt 0.0616 $X \sim B(250, "0.061551\dots")$ $Y \sim \text{Po}("15.3879\dots")$ [Accept $\text{Po}(15.4)$ or better] <u>or</u> suitable 3sf probs $= 0.14751\dots$ awrt <b>0.148</b>	B1	1.1b
		M1	3.3
		M1	3.4
		A1	1.1b
		(4)	
(c)	$H_0: \lambda = 16.8$ $H_1: \lambda < 16.8$ $U \sim \text{Po}(16.8)$ $P(U \geq 8) = 0.014$ [ $0.014 < 0.05$ or there is sufficient evidence to reject $H_0$ ] There is sufficient evidence at the 5% level of significance that the <u>number of calls received per day is lower in winter</u> <u>or rate of calls is lower in winter or less calls per day in winter</u> (o.e.)	B1	2.5
		B1	3.3
		M1	1.1b
		A1	2.2b
(d)	$C \sim \text{Po}(0.4 \times n + 0.2 \times n) [= \text{Po}(0.6n)]$ <u>or</u> $D \sim B(n, e^{-0.6})$ or awrt 0.549) $e^{-0.6n} < 0.001$ <u>or</u> $-0.6n < \ln(0.001)$ <u>or</u> $n > 11.5\dots$ $n = \underline{12}$	M1	3.1b
		M1	1.1b
		A1	1.1b
(e)	The <u>rate of calls per day is constant</u> <u>or</u> the <u>number of calls occurring in non-overlapping time intervals is independent.</u> <u>or</u> <u>number of calls per day is independent</u> (o.e.)	B1	2.4
		(1)	
<b>Total 14</b>			
(a)	<b>M1</b> <b>A1</b>	For using the model $\text{Po}(11.2)$ implied by sight of: 0.02077... or 0.9889.. or 0.9792.. awrt 0.021	
(b)	<b>B1</b>	awrt 0.0616	
	<b>1<sup>st</sup> M1</b>	Setting up a new model $B(250, "0.0616")$ [condone $B("0.0616", 250)$ ]	
	<b>2<sup>nd</sup> M1</b>	Seeing the model $\text{Po}(\text{their } np)$ implied by sight of: 0.1475.. or 0.89975 or 0.8524...	
SC	<b>A1</b>	awrt 0.148	
		if <b>no approximation</b> used (and 1 <sup>st</sup> M1 not seen) an answer of awrt 0.140 could get B1M1M0A0	
(c)	<b>1<sup>st</sup> B1</b>	Both hypotheses correct using $\lambda$ or $\mu$ and 16.8 or 0.4 [Accept their ans to $0.4 \times 42$ ]	
	<b>2<sup>nd</sup> B1</b>	Realising $\text{Po}(16.8)$ needs to be used. Sight or use of, implied by correct prob or CR	
	<b>M1</b>	For 0.014 or better (0.0141..) or CR $X \geq 9$ oe must be CR and not probability. [Allow CR $X \geq 10$ with probability $P(X \geq 10) = 0.054$ or better]	
	<b>A1</b>	<b>Indep of 1<sup>st</sup> B1</b> (must see 2 <sup>nd</sup> B1 and M1 scored) for a correct inference in context	
(d)	<b>1<sup>st</sup> M1</b>	Selecting a suitable model. Sight of $\text{Po}(0.6n)$ <u>or</u> $B(n, e^{-0.6})$ or implied by 2 <sup>nd</sup> M1	
	<b>2<sup>nd</sup> M1</b>	For a correct inequality or equality involving $n$ [Condone slips in solving] Allow <b>MR</b> i.e. misread of 0.01 for 0.001 (or similar) to score M1M1A0	
	<b>A1</b>	$n = 12$ cao [Correct answer with no incorrect working seen scores 3/3]	
(e)	<b>B1</b>	Allow equivalent statements. Underlined words required.	



Question	Scheme		Marks	AOs												
4(a) (i)	[ $W \sim \text{Geo}(0.11)$ ] $P(W = 6) = (0.89)^5 (0.11)$ $= 0.06142\dots$	awrt <b>0.0614</b>	M1	3.3												
			A1	1.1b												
(ii)	$P(W, 5) = 1 - (0.89)^5$ $= 0.44159\dots$	awrt <b>0.442</b>	M1	3.1b												
			A1	1.1b												
			(2)													
(iii)	$X \sim B(6, 0.11)$ $P(X = 4) = 0.001739\dots$	awrt <b>0.00174</b>	M1	3.3												
			A1	1.1b												
			(2)													
(iv)	[ $Y \sim \text{NB}(4, 0.11)$ ] using a neg bin $P(Y, 6) = P(Y = 4) + P(Y = 5) + P(Y = 6)$ $= (0.11)^4 + \binom{4}{3} (0.11)^3 (0.89)^1 \times 0.11 + \binom{5}{3} (0.11)^3 (0.89)^2 \times 0.11$ $= 0.001827$	or $V \sim B(6, 0.11)$ and $P(V \dots 4)$ for M2	M1	3.3												
			M1	3.1b												
			M1	3.4												
			A1	1.1b												
			(4)													
(b)	$P(\text{Zac wins}) = 0.89 \times 0.11 + (0.89)^3 \times 0.11 + (0.89)^5 \times 0.11 + \dots$ $= \frac{0.89 \times 0.11}{1 - (0.89)^2}$ oe $= 0.47089\dots = 0.471^*$		M1	3.1b												
			M1	1.1b												
			A1cso*	2.1												
			(3)													
<b>Total 13</b>																
(a)(i)	M1	Correct method to find $P(W = 6)$ eg $(p)^5 (1-p)$ for $p = 0.11$ or $0.89$														
	A1	awrt 0.0614 (Correct ans with no incorrect working 2/2)														
(ii)	M1	Correct method to find $P(W, 5)$														
	A1	awrt 0.442 (Correct ans with no incorrect working 2/2)														
(iii)	M1	For using the model $B(6, 0.11)$ allow $B(6, 0.89)$ [Implied by 0.0017 or awrt 0.114]														
	A1	awrt 0.00174 (Correct ans with no incorrect working 2/2)														
(iv)	1 <sup>st</sup> M1	In part (iv) we can accept correct expressions or values for probabilities For using a negative binomial model implied by correct $P(Y = 5)$ or $P(Y = 6)$														
	2 <sup>nd</sup> M1	Correct method to find $P(Y, 6)$														
	3 <sup>rd</sup> M1	At least two correct terms or $1 - 0.99817\dots$ from $1 - P(V, 3)$														
	A1	awrt 0.00183														
			<table border="1"> <thead> <tr> <th><math>a</math></th> <th>4</th> <th>5</th> <th>6</th> </tr> </thead> <tbody> <tr> <td><math>P(Y = a)</math></td> <td><math>1.46 \times 10^{-4}</math></td> <td><math>5.21 \times 10^{-4}</math></td> <td><math>1.16 \times 10^{-3}</math></td> </tr> <tr> <td><math>P(V = a)</math></td> <td><math>1.74 \times 10^{-3}</math></td> <td><math>8.60 \times 10^{-5}</math></td> <td><math>1.77 \times 10^{-6}</math></td> </tr> </tbody> </table>		$a$	4	5	6	$P(Y = a)$	$1.46 \times 10^{-4}$	$5.21 \times 10^{-4}$	$1.16 \times 10^{-3}$	$P(V = a)$	$1.74 \times 10^{-3}$	$8.60 \times 10^{-5}$	$1.77 \times 10^{-6}$
$a$	4	5	6													
$P(Y = a)$	$1.46 \times 10^{-4}$	$5.21 \times 10^{-4}$	$1.16 \times 10^{-3}$													
$P(V = a)$	$1.74 \times 10^{-3}$	$8.60 \times 10^{-5}$	$1.77 \times 10^{-6}$													
(b)	1 <sup>st</sup> M1	Forming the correct probability of Zac winning or identify $a$ and $r$ of GP	Allow for $p = (0.11) \times 0 + (1 - 0.11)(1 - p)$													
	2 <sup>nd</sup> M1	Using sum to infinity of a GP	Allow for $p = \frac{0.89}{1 + 0.89}$													
	A1*	Previous method marks must be seen leading to an answer 0.471 (NOT awrt 0.471)														

Question	Scheme	Marks	AOs
5	<p>Geo (0.3) <math>\mu = \frac{1}{0.3}</math> [ or exact equivalent e.g. <math>\frac{10}{3}</math> ]</p> <p><math>\sigma^2 = \frac{1-0.3}{0.3^2}</math> [ or exact equivalent e.g. <math>\frac{70}{9}</math> ]</p> <p>CLT <math>\Rightarrow \bar{X} \approx N\left(\frac{10}{3}, \dots\right)</math> oe</p> <p><math>\Rightarrow \bar{X} \approx N\left(\frac{10}{3}, \frac{7}{135}\right)</math> and attempt (sight of) <math>P(\bar{X} &lt; 3.45)</math></p> <p style="text-align: center;"><math>= 0.69579\dots</math> awrt <b>0.696</b></p>	B1 B1 M1 M1 A1	1.1b 1.1b 2.1 3.4 1.1b
<b>Total 5</b>			
<b>1<sup>st</sup> B1</b>	correct mean		
<b>2<sup>nd</sup> B1</b>	correct Var may be implied by sight of $\frac{7}{135}$ in distribution of $\bar{X}$		
<b>1<sup>st</sup> M1</b>	For use of CLT (must see $\bar{X}$ and Normal with mean correct ft) <u>or</u> sight of $N\left(\frac{10}{3}, \frac{7}{135}\right)$ <u>or</u> $N\left(\frac{10}{3}, \frac{70}{9 \times 150}\right)$ with any letter		
	Allow 3.33 or better for $\frac{10}{3}$ and 7.78 or better for $\frac{70}{9}$		
	May be implied by 2 <sup>nd</sup> M1		
<b>2<sup>nd</sup> M1</b>	Using the normal distribution to find $P(\bar{X} < 3.45)$ ft their " $\frac{10}{3}$ " and " $\frac{70}{150}$ "		
	May be implied by correct answer.		
<b>A1</b>	awrt 0.696		
Correct answer with no incorrect working scores 5/5			
	<b>Alternative (Use of <math>Y = \sum X</math>)</b>		
	$\mu = \frac{150}{0.3} [= 500]$	B1	
	$\sigma^2 = \frac{150 \times 0.7}{0.3^2} \left[ \frac{3500}{3} \right] = 1166.\dot{6}$	B1	
	$\Rightarrow Y \approx N\left(500, \frac{3500}{3}\right)$	M1	
	$P(Y < 517.5)$	M1	
	$= 0.69579\dots$	A1	

Question	Scheme	Marks	AOs
<b>6(a)</b>	$G_v(t) = \frac{9}{25}t^2 + \frac{12}{25}t^3 + \frac{4}{25}t^4$ <u>or</u> $t^2\left(\frac{9}{25} + \frac{12}{25}t + \frac{4}{25}t^2\right)$	M1	1.1b
	$= t^2\left(\frac{2}{5}t + \frac{3}{5}\right)^2$ *	A1* cso	2.1
		(2)	
<b>(b)(i)</b>	$G_w'(t) = 2t\left(\frac{2}{5}t + \frac{3}{5}\right)^4 + \left(\frac{2}{5}t + \frac{3}{5}\right)^5$	M1	2.1
	$[G_w'(1) = ]$ <u>3</u>	A1	1.1b
<b>(ii)</b>	$G_w''(t) = 2\left(\frac{2}{5}t + \frac{3}{5}\right)^4 + \frac{16}{5}t\left(\frac{2}{5}t + \frac{3}{5}\right)^3 + 2\left(\frac{2}{5}t + \frac{3}{5}\right)^4$ oe	M1	2.1
	$G_w''(1) = \frac{36}{5}$	A1	1.1b
		M1	2.1
		A1	1.1b
		(6)	
<b>(c)</b>	$G_x(t) = t^2\left(\frac{2}{5}t + \frac{3}{5}\right)^2 \times t\left(\frac{2}{5}t + \frac{3}{5}\right)^5$	M1	3.1a
	$= t^3\left(\frac{2}{5}t + \frac{3}{5}\right)^7$	A1	1.1b
		(2)	
<b>(d)</b>	$G_Y(t) = t^3 \times (t^2)^3 \times \left(\frac{2}{5}t^2 + \frac{3}{5}\right)^7$	M1	3.1a
	$= t^9\left(\frac{2}{5}t^2 + \frac{3}{5}\right)^7$	A1	1.1b
		(2)	
<b>(e)</b>	P(Y = 15) is coefficient of $t^{15}$ ie $\dots + t^9 \times {}^7C_3 \left(\frac{2}{5}t^2\right)^3 \left(\frac{3}{5}\right)^4 + \dots$	M1	1.1b
	<u>or</u> P(X = 6) need coefficient of $t^6$ i.e. $\dots + t^3 \times {}^7C_3 \left(\frac{2}{5}t\right)^3 \left(\frac{3}{5}\right)^4 + \dots$		
		A1	1.1b
		(2)	
<b>(14 marks)</b>			

Notes:		
(a)	M1 A1*	A correct un-simplified pgf based on $\sum t^v P(V = v)$ cso must see an un-simplified version i.e. M1 scored and no incorrect working seen
(b) (i)	M1 A1	Differentiating using the product rule to find $G_w'(t)$ Allow un-simplified e.g. $5 \times \frac{2}{5}t$ Need two terms added and at least one correct. If they expand we need 3 correct. 3 from a correct derivative
(ii)	1 <sup>st</sup> M1 1 <sup>st</sup> A1 2 <sup>nd</sup> M1 2 <sup>nd</sup> A1	Attempt $G_w''(t)$ ft their $G_w'(t)$ [must be at least 2 terms or a product], one correct ft term, same rule for differentiating a product $\frac{36}{5}$ or 7.2 from a correct derivative $G_w''(1) + G_w'(1) - (G_w'(1))^2$ ft their $G_w''(t)$ if different from $G_w'(t)$ and $G_w(t)$ <b>Dep on M3A2</b> $\frac{6}{5}$ or 1.2
(c)	M1 A1	Realising the need to use $G_x(t) = G_v(t) \times G_w(t)$ $t^3 \left( \frac{2}{5}t + \frac{3}{5} \right)^7$
(d)	M1 A1	Realising the need to multiply through by $t^3$ or substitute $t^2$ for $t$ or sight of $t^3 G_x(t^2)$ $t^9 \left( \frac{2}{5}t^2 + \frac{3}{5} \right)^7$ oe Need not be in its simplest form
(e)	M1 A1	Attempting to find correct coefficient of $t^n$ or identify $Y = 2J + 9$ where $J \sim B(7, 0.4)$ Need an expression can ft their $G_Y(t)$ or $G_X(t)$ of the form $t^n (at^m + b)^k$ Allow a statement that $P(Y = 15) = 0$ if it follows from their pgf For a correct exact answer or allow awrt 0.2903 Allow 0.29 from correct expression

#### Alternative for (b)

(b)	$W = P + 1$ where $P \sim B(5, 0.4)$ so $\text{Var}(W) = \text{Var}(P)$		
(i)	$G_P'(t) = 2 \left( \frac{2}{5}t + \frac{3}{5} \right)^4$	M1	2.1
	$G_w'(1) = 2 + 1 = 3$	A1	1.1b
(ii)	$G_P''(t) = \frac{16}{5} \left( \frac{2}{5}t + \frac{3}{5} \right)^3$ ; $G_P''(1) = \frac{16}{5}$	M1; A1	2.1 1.1b
	$\text{Var}(W) = \frac{16}{5} + 2 - (2)^2$ ; $= \frac{6}{5}$	M1; A1	2.1 1.1b
SC	<b>MR</b> They use $G_v(t)$ instead of $G_w(t)$ Provided some correct differentiation seen: Award B1 for $E(V) = \frac{14}{5}$ and B1 for $\text{Var}(V) = \frac{12}{25}$ score as M0A1M0A0M0A1		

Question	Scheme	Marks	Aos
7(a)	$\bar{X} \sim N(1000, 90)$ (May be implied by correct prob or $z$ value seen) $P(\bar{X} > 1020) = 0.0175\dots$ or $z = 2.108$ $0.0175\dots < 0.025$ or $z = 2.108\dots > 1.96$ therefore reject $H_0$ . There is evidence that the <u>mean weight</u> of the <u>flour</u> in a bag is <u>not 1000 g</u> <b>or</b> evidence of a <u>change</u> in <u>mean weight</u> of <u>flour</u> in a bag	M1 A1 M1 A1 cso <b>(4)</b>	3.3 3.4 1.1b 2.2b
(b)	$\left[ \bar{Y} \sim N\left(1000, \frac{900}{n}\right) \Rightarrow \frac{c-1000}{30/\sqrt{n}} = 1.6449 \right]$ $c = 1000 + \frac{49.347}{\sqrt{n}}$	M1 A1 <b>(2)</b>	3.4 1.1b
(c)	$\frac{1000 + \frac{49.347}{\sqrt{n}} - 1020}{30/\sqrt{n}} = -2.5758$ $\frac{126.621}{\sqrt{n}} = 20 \quad \text{or} \quad \frac{49.34\dots}{c-1000} = \frac{-77.274}{c-1020} \quad (\text{Allow 2sf accuracy})$ $n = \underline{40}$ $c = 1007.8\dots$ <p style="text-align: right;">awrt <b><u>1010</u></b></p>	M1 A1ft dM1 A1 A1 <b>(5)</b>	3.4 1.1b 1.1b 2.1 1.1b

**(11 marks)**

**Notes:**

(a)	<b>1<sup>st</sup> M1</b> Setting up the correct model. Normal with $\mu = 1000, \sigma^2 = 90$ or $\sigma = \sqrt{90}$ or awrt 9.49 <b>1<sup>st</sup> A1</b> Using the model to find the correct $z$ value or $P(\bar{X} > 1020) =$ awrt 0.0175 Allow CR $\bar{C} \dots 1018.59\dots$ awrt 1019 [ $>$ is OK] Ignore lower CR provided $< 1000$ <b>2<sup>nd</sup> M1</b> Correct comparison <u>or</u> non-contextual conclusion. Allow comparison of 1020 with critical region. <b>Dep on</b> $P(\bar{X} > 1020)$ M0 if there are contradictory statements. <b>2<sup>nd</sup> A1</b> cso <b>dep on M1A1M1</b> for a correct conclusion in context with underlined words Do NOT accept “mean weight has <u>increased</u> ”
(b)	<b>M1</b> For Finding the CR using the Normal distribution. Condone $\sigma = \sqrt{\frac{30}{n}}$ to score M1 $\frac{c-1000}{30/\sqrt{n}} = z$ where $ z  > 1.5$ <b>Allow any inequality or = for M1 in (b) and M1 A1ft M1 in (c)</b> <b>A1</b> A correct equation in the form $c = \dots$ and for use of awrt 1.6449 (implied by awrt 49.3[4]) Condone $\bar{X}$ used for $c$ (o.e.)
(c)	<b>1<sup>st</sup> M1</b> Standardising using their $c$ (letter or expression) and equating to $z$ ( $ z  > 2$ ) to form an equation in $n$ or $n$ and $c$ . Can ft their $\sigma$ <b>used</b> in (b) for M1A1ft here <b>1<sup>st</sup> A1ft</b> Ft their “ $c$ ” for a correct equation with $-2.58$ (or 1.64 or 1.65 used in (b)) <b>2<sup>nd</sup> dM1</b> <b>Dependent 1<sup>st</sup> M1</b> . Isolating or eliminating either $\sqrt{n}$ or $n$ <u>or</u> eliminating $c$ leading to an equation for $n$ or $c$ <b>2<sup>nd</sup> A1</b> For 40 (allow 41) Must be an integer. With correct working.   e.g. Check correct $\sigma$ <b>3<sup>rd</sup> A1</b> For awrt 1010 from correct working   has been used